

**WARNING:**  
READ LABEL BEFORE USE  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

# BANTI<sup>®</sup> POUR-ON

FOR THE CONTROL OF TICKS ON CATTLE AND  
AS AN AID IN THE TREATMENT OF TICKS ON DEER

Active ingredient: 10 g/L flumethrin

NET CONTENTS: 5L

PLBAS5F-01



# BANTIX<sup>®</sup> POUR-ON

## FOR ANIMAL TREATMENT ONLY

BANTIX<sup>®</sup> Pour-On contains 10 g/L flumethrin. It is active against all 3 stages of the New Zealand cattle tick (*Haemaphysalis longicornis*) lifecycle. It also causes infertility of ticks which survive treatment, by inhibiting egg laying or rendering eggs sterile. This action helps reduce contamination of pasture.

### DIRECTIONS FOR USE.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Apply BANTIX<sup>®</sup> Pour-On evenly along the midline of the back from the front of the shoulders to the tails using a pour-on applicator. Check accuracy of pour-on applicator regularly. BANTIX<sup>®</sup> Pour-On spreads through the oils of the skin and hair coat of all parts of the animal's body. Behaviour such as grooming, tail swishing and scratching, aid the movement of active ingredient to the extremities of the body, which are often affected by ticks. Retreatment should be under taken when ticks start to reappear.

For optimum control in deer, breeding hinds should be treated every three weeks from August to fawning. This program will reduce the number of nymphs reaching the adult stage, which infest fawn during November/December. Treatment of hinds can be continued after fawning if this is practical. Other stock should be treated every three weeks from August, or as required.

### Cattle: Dose Table

Live Weight	Dose Volume
Up to 200kg	20ml
200 - 300kg	30ml
300 - 400kg	40ml
> 400kg	10ml/100kg

### Deer: Dose Table

Age	Dose Volume
Up to 6 months	10ml
Yearlings & adults	20ml

**HANDLING PRECAUTIONS:** May cause a temporary skin irritation in some individuals if applied to the skin. Avoid contact with hands and skin. Wear protective clothing and gloves when handling. Do not breath vapour.

**First aid:** If on skin wash off with soap and water. If you feel unwell get medical advice. Immediately call a POISON CENTER (0800 764 766) or doctor.

**ENVIRONMENTAL PROTECTION:** Harmful to aquatic life with long lasting effects. Avoid contamination of any water supply with product or empty container. Harmful to terrestrial vertebrates. Avoid release into the environment.

**DISPOSAL:** Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved facility. Triple rinse empty container. Recycle empty container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved facility. Do not use the container for any other purpose.

### ADVERSE EFFECTS:

Signs of mild irritation may be observed for 24-48 hours after treatment. Hair loss at the site of application may occur on deer which are shedding their winter coat at the time of treatment. This effect is temporary and can be minimized by spreading the dose over the entire backline. Growth of summer coat is not affected.

### WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

Meat: Nil

Milk: Nil

EPA Approval No. HSR100759

See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

ACVM Registration No. A11154

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

STORE below 25°C. Do not freeze.


PLBA5B-01

B.  
EXP.

FREE RECYCLING



[agrecovery.co.nz](http://agrecovery.co.nz)



0 799439 606258



Registered to  
Alleva Animal Health Ltd  
15 Calman Place,  
Birkenhead, Auckland