

WARNING
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

CORPORAL[®]

MINERALISED DRENCH FOR SHEEP

A BROAD SPECTRUM COMBINATION SHEEP DRENCH FOR THE CONTROL OF SUSCEPTIBLE GASTROINTESTINAL ROUNDWORMS AND LUNGWORMS INCLUDING BENZIMIDAZOLE AND LEVAMISOLE RESISTANT STRAINS.

Ingredients: Albendazole 23.8mg/mL, Levamisole HCl 37.5mg/mL, Selenium 0.5mg/mL (as sodium selenate 1.2mg/mL), Cobalt 0.3mg/mL, Copper 2.1mg/mL, Zinc 0.6mg/mL

NET CONTENTS: 5L

PLCMS5F-01



CORPORAL[®]
MINERALISED DRENCH FOR SHEEP

PLEASE READ LABEL
COMPLETELY BEFORE USE

INDICATIONS: Contains albendazole 23.8mg/mL, levamisole HCl 37.5mg/mL, selenium 0.5mg/mL, cobalt 0.3mg/mL, copper 2.1mg/mL, zinc 0.6mg/mL. For the treatment of tapeworms and liver fluke and the following susceptible immature and mature gastrointestinal roundworms and lungworms, including benzimidazole and levamisole resistant strains: *Haemonchus contortus*, *Ostertagia* spp., *Trichostrongylus* spp., *Nematodirus* spp., *Cooperia* spp., *Strongyloides papillosus*, *Bunostomum trigonocephalum*, *Chapertia ovina*, *Oesophagostomum venulosum*, *Trichuris ovis* (adult only), *Moniezia expansa* (segments only), *Dictyocaulus* spp., *Muellerius capillaris*, Liver fluke (adults only). Also, reduces the output of roundworm and fluke eggs.

WARNING: DANGEROUS TO THE ENVIRONMENT. VERY TOXIC TO AQUATIC ORGANISMS. AVOID CONTAMINATION OF ANY WATER SUPPLY WITH PRODUCT OR EMPTY CONTAINER. HARMFUL TO THE SOIL ENVIRONMENT.

RESISTANCE MANAGEMENT: Resistance can develop to any anthelmintic. Ask your local animal health advisor or veterinary practitioner for recommended parasite management practices for your area to reduce the development of resistance. It is advised that a resistance test be carried out regularly when using any parasite treatment.

DIRECTIONS FOR USE: By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Shake well before use. CORPORAL Mineralised Drench for Sheep is a ready to use anthelmintic that can be dosed through standard drenching equipment. It is important to give the correct dose. Therefore, before treatment, weigh a representative sample of sheep to determine the weight of the heaviest animal. Where there is a large variation in liveweights, drafting into two or more lines may be appropriate before weighing and treating, to avoid under and over dosing. In this situation it is important to choose the dose rate based on the label recommendation for each weight band. Use correct drenching techniques and check accuracy of drenching equipment regularly. Wash out drench gun thoroughly with water before and after use.

DOSE CHART: Dose rate sheep: 1mL/5kg
(4.75mg/kg albendazole, 7.5mg/kg levamisole HCL)

Animal weight	Dose	No. of doses per 5L pack
15kg	3 mL	1666
16-20kg	4 mL	1250
21-25kg	5 mL	1000
26-30kg	6 mL	833
31-35kg	7 mL	714
36-40kg	8 mL	625
41-45kg	9 mL	555
46-50kg	10 mL	500
51-55kg	11 mL	454
56-60kg	12 mL	416
61-65kg	13 mL	384
66-70kg	14 mL	357

Over 70kg dose at the rate of 1mL/5kg

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

WITHHOLDING PERIODS:

MEAT: Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 10 days of the last treatment

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment

STORAGE: Store locked up in original container, tightly closed, in cool, dry conditions. Protect from light. Avoid temperatures below 0° and above 30°C.

Registered Pursuant to the ACVM Act 1997, No. A11398
See www.foodsafety.govt.nz/acvm for registration conditions.


Batch No.

Expiry Date.



Registered to
Alleva Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland

HANDLING PRECAUTIONS: May be harmful if swallowed. Repeated exposure may cause skin allergy and mild skin irritation. Albendazole and levamisole hydrochloride may possibly cause damage to genetic material and may affect development and/or reproduction. Levamisole hydrochloride may possibly affect blood and haemopoietic system. Avoid skin contact. Wear gloves when decanting. Do not eat, drink or smoke when using this product. Avoid release into the environment. **PRECAUTION:** Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian. Do not exceed stated dose or dose more frequently than three weekly. In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper poisoning (deaths) occurring following copper supplementation in this species. Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity so estimate liveweights carefully. Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between levamisole and organophosphate dips. **FIRST AID:** If swallowed, Do NOT induce vomiting. For advice, contact a doctor or the National Poisons Centre 0800 POISON (0800 764 766) immediately. If skin or hair contact occurs, remove contaminated clothing and wash skin thoroughly. If splashed in eyes, flush immediately with copious amounts of water. **EMERGENCY RESPONSE:** In case of spillage, wear appropriate protective clothing and prevent material from entering water ways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely. **DISPOSAL:** Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.



FREE RECYCLING



agrecovery.co.nz



PLCMS5B-01