

WARNING:
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

CORPORAL^{+TAPE}

MINERALISED DRENCH FOR SHEEP

A BROAD SPECTRUM SHEEP DRENCH FOR THE CONTROL OF TAPEWORMS (HEADS AND SEGMENTS), ADULT LIVER FLUKE AND SUSCEPTIBLE GASTROINTESTINAL ROUNDWORMS AND LUNGWORMS IN SHEEP AND LAMBS INCLUDING BENZIMIDAZOLE AND LEVAMISOLE RESISTANT STRAINS.

Ingredients: Albendazole 25mg/mL, Levamisole HCl 37.5mg/mL, Praziquantel 18.8mg/mL, Selenium 0.5mg/mL, Cobalt 0.2mg/mL, Copper 2mg/mL, Zinc 0.6mg/mL, Iodine 1mg/mL

NET CONTENTS: 5L

PLCORTMS5F-01



CORPORAL⁺TAPE

MINERALISED DRENCH FOR SHEEP

PLEASE READ LABEL COMPLETELY BEFORE USE

INDICATIONS: Contains albendazole 25mg/mL, levamisole HCl 37.5mg/mL praziquantel 18.8mg/mL, selenium 0.5mg/mL, cobalt 0.2mg/mL, copper 2mg/mL, zinc 0.6mg/mL, iodine 1mg/mL. For the treatment of tapeworm (*Moniezia spp.*) heads and segments in lambs and sheep. For the treatment of the following susceptible immature and mature [*includes inhibited L4] gastrointestinal roundworms and lungworms, including benzimidazole and levamisole resistant strains: *Haemonchus spp.* *, *Ostertagia spp.* *, *Cooperia spp.* *, *Trichostrongylus spp.*, *Oesophagostomum spp.* [adults only] *Nematodirus spp.* [adults only], *Dictyocaulus spp.* For the treatment of adult liver fluke.

RESISTANCE MANAGEMENT: Resistance can develop to any anthelmintic. Ask your local animal health advisor or veterinary practitioner for recommended parasite management practices for your area to reduce the development of resistance. It is advised that a resistance test be carried out regularly when using any parasite treatment.

WARNING: DANGEROUS TO THE ENVIRONMENT. VERY TOXIC TO AQUATIC ORGANISMS AND HARMFUL TO THE SOIL ENVIRONMENT. AVOID RELEASE INTO THE ENVIRONMENT AND CONTAMINATION OF ANY WATER SUPPLY WITH PRODUCT OR EMPTY CONTAINER

DIRECTIONS FOR USE: By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Shake well before use. CORPORAL + Tape Mineralised Drench for Sheep is a ready to use anthelmintic that can be dosed through standard drenching equipment. It is important to give the correct dose. Therefore, before treatment, weigh a representative sample of sheep to determine the weight of the heaviest animal. Where there is a large variation in liveweights, drafting into two or more lines may be appropriate before weighing and treating, to avoid under and over dosing. In this situation it is important to choose the dose rate based on the label recommendation for each weight band.

Use correct drenching techniques and check accuracy of drenching equipment regularly. Wash out drench gun thoroughly with water before and after use.

DOSE CHART: Dose rate sheep: 1mL/5kg (5mg/kg albendazole, 7.5mg/kg levamisole HCL and 3.76g/kg praziquantel)

Liveweight	Dose	5L pack treats
15kg	3mL	1666
16kg - 20kg	4mL	1250
21kg - 25kg	5mL	1000
26kg - 30kg	6mL	833
31kg - 35kg	7mL	714
36kg - 40kg	8mL	625
41kg - 45kg	9mL	555
46kg - 50kg	10mL	500
51kg - 55kg	11mL	454
56kg - 60kg	12mL	416
61kg - 65kg	13mL	384
66kg - 70kg	14mL	357

Over 70kg dose at the rate of 1mL/5kg

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Limits of Agricultural Compounds.

MEAT: Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 10 days of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.



Storage: Store in original container, tightly closed, in cool, dry conditions. Protect from light. Avoid temperatures below 0° and above 25°C.

Registered Pursuant to the ACVM Act 1997, No. A11351
See www.foodsafety.govt.nz/acvm for registration conditions

B.

EXP.

HANDLING PRECAUTIONS: May be harmful if swallowed. Handle with care avoiding skin contact. May cause mild skin irritation. Repeated exposure may cause skin allergy. Albendazole and levamisole hydrochloride may possibly cause damage to genetic material and may affect development and/or reproduction. Levamisole hydrochloride may possibly affect blood and haemopoietic system. **PRECAUTIONS:** Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian. Do not exceed stated dose or dose more frequently than three weekly. In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper poisoning (deaths) occurring following copper supplementation in this species. Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity so estimate liveweights carefully. Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between levamisole and organophosphate dips. **FIRST AID:** If swallowed, Do NOT induce vomiting. For advice, contact a doctor or the National Poisons Centre 0800 POISON (0800 764 766) immediately. If skin or hair contact occurs, remove contaminated clothing and wash skin thoroughly. If splashed in eyes, flush immediately with copious amounts of water. **DISPOSAL:** Preferably dispose of product by use, otherwise dispose of product and packaging at an approved landfill or other approved facility. **EMERGENCY RESPONSE:** In case of spillage, wear appropriate protective clothing and prevent material from entering water ways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely.



Registered to
Alleva Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland

FREE RECYCLING



agrecovery.co.nz

PLCORTMS5B-01