

WARNING:
READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

REFLEX[®]

POUR-ON

FOR THE TREATMENT AND CONTROL OF
INTERNAL AND EXTERNAL PARASITES OF CATTLE

Active ingredient: 10 g/L ABAMECTIN
NET CONTENTS: 5L

PLRP5F-01



REFLEX® POUR-ON

FOR ANIMAL TREATMENT ONLY

REFLEX Pour-on contains 10mg/ml abamectin, a member of the macrocyclic lactone group of compounds. It is effective against internal and external parasites sensitive to this family. REFLEX Pour-on when used at the recommended dose level of 0.5mg abamectin per kg liveweight effectively controls the following abamectin-sensitive parasites of cattle:

Gastro-intestinal roundworms: Adult and immature Small Brown Stomach Worm* (*Ostertagia spp.*), Barber's Pole Worm (*Haemonchus spp.*), Stomach Hair Worm and Black Scour Worm (*Trichostrongylus spp.*), Small Intestinal Worm (*Cooperia spp.*), Hookworm (*Bunostomum spp.*), Large Bowel Worm (*Oesophagostomum spp.*), and adults of Large-Mouthed Bowel Worm (*Chabertia spp.*), Thin-Necked Intestinal Worm (*Nematodirus spp.*), Whipworm (*Trichuris spp.*) and Intestinal Threadworm (*Strongyloides spp.*) *including inhibited immature

Lungworms: Adult and immature *Dictyocaulus viviparus*.

Biting and Sucking Lice: *Bovicola bovis*, *Linognathus vituli*.

Resistance can develop to any chemical. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

Production Enhancement: Increases milk production in grazing dairy cows. Data has shown that milk production increases by a median of +0.63kg liquid milk per day when cows or heifers infected with gastrointestinal nematode parasites are treated with an effective anthelmintic; this increase occurs without any reduction in milk quality.

DIRECTIONS FOR USE.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Ready for use with standard pour-on equipment. Check accuracy of pour-on applicator regularly.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Apply along the backline from shoulders to tail.

DO NOT treat cattle that weigh less than 100kg

DOSE: 1ml per 20kg bodyweight
(0.5 mg abamectin per kg bodyweight)

Liveweight	Dose	5L pack treats
50-100kg	5 mL	1000
101-150kg	7.5 mL	666
151-200kg	10 mL	500
201-250kg	12.5 mL	400
251-300kg	15 mL	333
301-350kg	17.5 mL	285
351-400kg	20 mL	250
401-450kg	22.5 mL	222
451-500kg	25 mL	200
501-550kg	27.5 ml	181
551-600kg	30 ml	166
601-650kg	32.5ml	153

Cattle in excess of 650 kg to be dosed at 1 mL per 20 kg bodyweight.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

MEAT: Cattle producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 35 days of last treatment.

MILK: Nil.

EPA Approval No. HSR100758

See www.epa.govt.nz for approval conditions

ACVM Registration No. A10666

See www.foodsafety.govt.nz for registration conditions

STORE below 25°C B.

EXP.

Read label and obtain special instructions before use.

WARNING: Harmful if swallowed, inhaled or in contact with skin. May cause harm to breast-fed children. Very toxic to aquatic organisms. Harmful to soil environment, terrestrial vertebrates and invertebrates.

HANDLING PRECAUTIONS: Wear protective gloves and clothing. Use only outdoors or in a well ventilated area. Avoid breathing vapour. Avoid contact during pregnancy/while nursing. Do not eat, drink or smoke when using this product. Wash hands thoroughly after handling. Wash contaminated clothing and/or protective equipment before reuse.

FIRST AID: IF SWALLOWED: contact the National Poisons Centre (0800 764 766) or a doctor immediately. IF ON SKIN: wash with plenty of soap and water. IF INHALED: remove to fresh air and keep at rest in a position comfortable for breathing. If exposed or concerned get medical advice immediately. If medical advice is need, have product container or label at hand.

DISPOSAL: Triple rinse empty container, crush or puncture, and bury in a suitable landfill, or if appropriate, recycle. Avoid contamination of any water supply with chemical or empty container.



FREE RECYCLING



agrecovery.co.nz



Registered to
Alleva Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland

WARNING:
READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

REFLEX[®]

POUR-ON

FOR THE TREATMENT AND CONTROL OF
INTERNAL AND EXTERNAL PARASITES OF CATTLE

Active ingredient: 10 g/L ABAMECTIN

NET CONTENTS: 20L



REFLEX[®] POUR-ON

FOR ANIMAL TREATMENT ONLY

REFLEX Pour-on contains 10mg/ml abamectin, a member of the macrocyclic lactone group of compounds. It is effective against internal and external parasites sensitive to this family. REFLEX Pour-on when used at the recommended dose level of 0.5mg abamectin per kg liveweight effectively controls the following abamectin-sensitive parasites of cattle:

Gastro-intestinal roundworms: Adult and immature Small Brown Stomach Worm* (*Ostertagia spp.*), Barber's Pole Worm (*Haemonchus spp.*), Stomach Hair Worm and Black Scour Worm (*Trichostrongylus spp.*), Small Intestinal Worm (*Cooperia spp.*), Hookworm (*Bunostomum spp.*), Large Bowel Worm (*Oesophagostomum spp.*), and adults of Large-Mouthed Bowel Worm (*Chabertia spp.*), Thin-Necked Intestinal Worm (*Nematodirus spp.*), Whipworm (*Trichuris spp.*) and Intestinal Threadworm (*Strongyloides spp.*)
*including inhibited immature

Lungworms: Adult and immature *Dictyocaulus viviparus*.

Biting and Sucking Lice: *Bovicola bovis*, *Linognathus vituli*.

Resistance can develop to any chemical. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

Production Enhancement: Increases milk production in grazing dairy cows. Data has shown that milk production increases by a median of +0.63kg liquid milk per day when cows or heifers infected with gastrointestinal nematode parasites are treated with an effective anthelmintic; this increase occurs without any reduction in milk quality.

DIRECTIONS FOR USE.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Ready for use with standard pour-on equipment. Check accuracy of pour-on applicator regularly. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Apply along the backline from shoulders to tail. **DO NOT** treat cattle that weigh less than 100kg

DOSE: 1ml per 20kg bodyweight (0.5 mg abamectin per kg bodyweight)

Liveweight	Dose	20L pack treats
101-150kg	7.5 mL	2664
151-200kg	10 mL	2000
201-250kg	12.5 mL	1600
251-300kg	15 mL	1332
301-350kg	17.5 mL	1140
351-400kg	20 mL	1000
401-450kg	22.5 mL	888
451-500kg	25 mL	800
501-550kg	27.5 mL	724
551-600kg	30 mL	664
601-650kg	32.5 mL	612

Cattle in excess of 650 kg to be dosed at 1 mL per 20 kg bodyweight.

Read label and obtain special instructions before use.

WARNING: Harmful if swallowed, inhaled or in contact with skin. May cause harm to breast-fed children.

HANDLING PRECAUTIONS: Wear protective gloves and clothing. Use only outdoors or in a well ventilated area. Avoid breathing vapour. Avoid contact during pregnancy/while nursing. Do not eat, drink or smoke when using this product. Wash hands thoroughly after handling. Wash contaminated clothing and/or protective equipment before reuse.

FIRST AID: IF SWALLOWED: contact the National Poisons Centre (0800 764 766) or a doctor immediately. IF ON SKIN: wash with plenty of soap and water. IF INHALED: remove to fresh air and keep at rest in a position comfortable for breathing. If exposed or concerned get medical advice immediately. If medical advice is needed, have product container or label at hand.

DISPOSAL: Triple rinse empty container, crush or puncture, and bury in a suitable landfill, or if appropriate, recycle. Avoid contamination of any water supply with chemical or empty container.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

MEAT: Cattle producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 35 days of last treatment.

MILK: Nil.

EPA Approval No. HSR100682

See www.epa.govt.nz for approval conditions

ACVM Registration No. A10666

See www.foodsafety.govt.nz for registration conditions

STORE below 25°C

B.

EXP.



Registered to
Alleva Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland



REFLEX[®] is a registered trademark of Alleva Animal Health Ltd